

1410 N. Hilton
Boise, ID 83706

DEPARTMENT OF ENVIRONMENTAL QUALITY

www.deq.idaho.gov



2007 Air Quality Monitoring Data Summary

Table of Contents

TABLE OF CONTENTS	III
APPENDIX—DATA TABLES	IV
LIST OF TABLES	V
LIST OF GRAPHS AND MAPS	VI
INTRODUCTION	1
AIR QUALITY STANDARDS	3
MONITORING NETWORK	5
PARTICULATE MONITORING	5
MONITORING RESULTS	9
OZONE	9
PARTICULATE MATTER (10 MICROMETERS)	15
PARTICULATE MATTER (2.5 MICROMETERS)	17
CARBON MONOXIDE	27
SULFUR DIOXIDE	29
LEAD	33
NITROGEN DIOXIDE	35
AIR QUALITY INDEX	37
IMPAIRED AIR QUALITY	49
WINTER BURN BANS	49
SUMMER OZONE ALERTS	49
DEFINITIONS	50
GENERAL DEFINITIONS	50
POLLUTION SOURCES	51
CRITERIA AIR POLLUTANTS	52
APPENDIX	A

Appendix—Data Tables

Air Quality Index

Calculation and Breakpoints for the Air Quality Index.....	A-2
EPA AirData Air Quality Index Report	A-3

2007 Data Tables

EPA AirData Monitor Values Report PM2.5.....	A-4
EPA AirData Monitor Values Report PM10.....	A-5
EPA AirData Monitor Values Report Ozone.....	A-6
EPA AirData Monitor Values Report CO.....	A-7
EPA AirData Monitor Values Report NO2.....	A-8
EPA AirData Monitor Values Report SO2	A-9
EPA AirData Monitor Values Report Lead	A-10

List of Tables

Air Quality Standards for Criteria Pollutants	3
Monitoring Network	6
Monitoring Methods Used in Idaho	7
Air Quality Index	37
Air Quality Index Ratings For Idaho Counties.....	48

List of Graphs and Maps

Air Monitoring Network	8
Lancaster 8-Hour Ozone	10
St. Luke’s 8-Hour Ozone.....	11
ITD 8-Hour Ozone	12
Whitney 8-Hour Ozone.....	13
Idaho 8-Hour Ozone Averages	14
Daily Maximum PM₁₀ 3-Year Average.....	16
Daily Maximum 98th% PM_{2.5} 3-Year Average.....	19
PM_{2.5} 3-Year Average Annual Mean	20
Boise PM_{2.5} Daily Averages from Continuous Analyzer	21
Nampa PM_{2.5} Daily Averages from Continuous Analyzer	21
Coeur d’Alene PM_{2.5} Daily Averages from Continuous Analyzer	22
St. Maries PM_{2.5} Daily Averages from Continuous Analyzer	22
Sandpoint PM_{2.5} Daily Averages from Continuous Analyzer	23
Pinehurst PM_{2.5} Daily Averages from Continuous Analyzer.....	23
Lewiston PM_{2.5} Daily Averages from Continuous Analyzer.....	24
Moscow PM_{2.5} Daily Averages from Continuous Analyzer	24
Twin Falls PM_{2.5} Daily Averages from Continuous Analyzer	25
Pocatello PM_{2.5} Daily Averages from Continuous Analyzer.....	25
Idaho Falls PM_{2.5} Daily Averages from Continuous Analyzer	26
Salmon PM_{2.5} Daily Averages from Continuous Analyzer	26
Carbon Monoxide (CO) 2nd Highest 8-Hour Concentration	28
Sulfur Dioxide (SO₂) Annual Average	30
Sulfur Dioxide (SO₂) Maximum 24-Hour Average.....	31
Sulfur Dioxide (SO₂) Maximum 3-Hour Average.....	32
Lead Maximum Quarterly Average.....	34
Nitrogen Dioxide Annual 1-Hour Average.....	36
Percentage of Days Air Quality Was Rated As “Good” per AQI	38
Air Quality for Ada County	39
Air Quality for Bannock County	40
Air Quality for Bonneville County	41
Air Quality for Canyon County.....	42
Air Quality for Franklin County	43
Air Quality for Kootenai County.....	44
Air Quality for Nez Perce County	45
Air Quality for Shoshone County.....	46
Air Quality for Twin Falls County.....	47

The 2007 Air Quality Monitoring Data Summary is available
for viewing or downloading on the internet at:

<http://www.deq.state.id.us/air/>

Links to additional documents for download are also available at the Web site.



2007 Air Quality Data Summary

Introduction

This annual report is issued by the Idaho Department of Environmental Quality (DEQ) to inform the public of air quality throughout Idaho. The purpose of this report is to summarize regional ambient air quality while presenting air monitoring results for six criteria air pollutants. The U.S. Environmental Protection Agency (EPA) sets national ambient air quality standards (NAAQS) for these pollutants. These criteria air pollutants are:

- Particulate Matter ($PM_{10} \leq 10$ micrometers, $PM_{2.5} \leq 2.5$ micrometers in diameter)
- Carbon Monoxide
- Sulfur Dioxide
- Nitrogen Dioxide
- Ozone
- Lead

In Idaho, monitoring for the criteria pollutants occurs primarily in areas of high population where the potential for human exposure is greatest. Particulate matter is currently the most common criteria pollutant of concern in Idaho because particulate sources are widespread throughout the state. Common sources include windblown dust, re-entrained road dust, smoke (residential, agricultural, and forest fires), industrial emissions, and motor vehicle emissions. The PM_{10} standard has been in effect since 1987 and historically has been the particulate size of concern. However, $PM_{2.5}$, or PM Fine, has been monitored in Idaho since 1998 and is now the pollutant of concern. Numerous studies have associated fine particulate matter with a variety of respiratory and cardiovascular problems, ranging from aggravated asthma, to irregular heartbeats, heart attacks, and early death in people with heart or lung disease. The $PM_{2.5}$ and PM_{10} NAAQS were changed by EPA effective December 17, 2006. Due to a lack of evidence linking health problems to long-term exposure to coarse particle pollution, EPA revoked the annual PM_{10} standard. The 24 hour standard for $PM_{2.5}$ was lowered from $65 \mu\text{g}/\text{m}^3$ to $35 \mu\text{g}/\text{m}^3$ to provide increased protection against health effects associated with short-term exposure (including premature mortality and increased hospital admissions and emergency room visits).

Another historical pollutant of concern in Idaho is carbon monoxide (CO). The primary source of CO is incomplete fossil fuel combustion. CO concentrations have the potential to be high in the urbanized areas where automobile traffic is heavy and cars frequently idle at stoplights. The Boise area (Northern Ada County) was the only CO nonattainment area in the state. When the State Implementation Plan (SIP) and Maintenance Plan were accepted by EPA, it was reclassified as a maintenance area on December 27, 2002. No violations of the 1- or 8-hour CO NAAQS have occurred since 1991.

Sulfur dioxide (SO_2) and nitrogen dioxide (NO_2) sources are few and localized because these pollutants come primarily from large industrial sources (transportation sources also contribute to NO_2). There is little heavy industry in Idaho and elevated SO_2 and NO_2 concentrations in ambient air are typically not found. However, due to concerns of some localized sources, DEQ has monitored for one or both of these pollutants in Boise, Pocatello, Moyie Springs, Mountain Home, and Soda Springs. In the past 10 years of targeted monitoring, DEQ has not measured significant concentrations of these pollutants at these



2007 Air Quality Data Summary

monitoring sites. DEQ initiated NO₂ monitoring near Coeur d'Alene on January 1, 2005 to characterize emissions in the area.

The fifth criteria pollutant monitored by DEQ is ozone. Ozone (O₃) has been monitored in the Treasure Valley since 2002, and in Coeur d'Alene beginning in 2005. Ozone is created when combustion by-products (volatile organic compounds, VOC) near the ground react with nitrogen oxides and other compounds to create photochemical smog. These reactions are stimulated on days of intense sunlight and warm temperatures. Ozone has become a pollutant of concern since many summertime days are classified as moderate for ozone on the Air Quality Index (AQI). EPA lowered the ozone standard, effective May 27, 2008. The new 8-hour ozone NAAQS is 0.075 ppm, lowered from 0.08 ppm. This new standard poses a greater risk of non-attainment for all airsheds, but particularly the Treasure Valley airshed.

Lead (Pb) is the sixth criteria pollutant and is not currently being monitored by DEQ. Lead (Pb) was monitored in the Shoshone County town of Kellogg, near the Bunker Hill superfund site, because lead was a by-product of the smelting process that occurred in the area for decades. Although a significant problem in the 1970s and early 1980s, airborne Pb concentrations at this monitoring site were very low through the 1990s. DEQ discontinued monitoring for lead in 2002. EPA is currently reviewing the lead NAAQS and is proposing to lower the standard significantly. Whether this lowering of the Pb NAAQS will trigger new monitoring in Idaho will not be determined until EPA decides on the NAAQS and the rule is final.

DEQ monitored for toxic air pollutants in the Treasure Valley from 2003 to the beginning of 2005 to determine if concentrations are at levels that could have adverse health effects. These health effects include, but are not limited to, increased cancer risk and respiratory, cardiovascular, and neurological effects. EPA, through their National Air Toxics Assessment (NATA) program, predicts cancer and non-cancer risk values across Idaho's airsheds using emissions estimates of certain air toxic compounds.

While Idaho generally enjoys good air quality, in many ways our airsheds are faced with new challenges. Some of these challenges are related to economic and population growth, particularly in terms of vehicles on roadways and growth in new construction.

Each day, DEQ measures the concentration of certain air pollutants throughout the state. Based on local ordinances, DEQ may issue a burn ban when concentrations of these air pollutants reach or exceed the health-based standards or limits established by state law or regulation. Concerned citizens may tune in to the news on their local radio or television station to find out if a burn ban has been issued, or login to DEQ's Internet web site at www.deq.state.id.us/air/index.cfm. DEQ issues a news bulletin to local news media, law enforcement, and fire officials each time a burn ban is imposed. There were a number of voluntary and mandatory bans issued in 2007.

Real-time air monitoring data are available on the Internet at www.tcsn.net/family/Idaho/index.html. We encourage you to visit our Web site at <http://www.deq.state.id.us/> to find more extensive air quality data, educational materials, and discussions of current topics.



2007 Air Quality Data Summary

We are expanding and refining our Internet site to better serve the residents of Idaho. We want your feedback on our air quality data and program. Please submit your comments via email to Bruce Louks; Monitoring, Modeling, & Emissions Inventory Manager, at Bruce.Louks@deq.idaho.gov or call at 208-373-0294.

Air Quality Standards

The federal Clean Air Act (CAA) requires EPA to set [National Ambient Air Quality Standards](#) (NAAQS) for pollutants considered harmful to public health and the environment. The standards are designed to primarily protect the general public, including sensitive populations such as asthmatics, children, and the elderly. They are also intended to safeguard public welfare by reducing effects such as decreased visibility and damage to animals, crops, vegetation, and buildings. EPA has established standards for six criteria pollutants. The list below contains seven pollutants, which include two size ranges of particulate matter.

The state of Idaho adheres to these standards. For more information, EPA air quality standards and supporting rationale are available at <http://epa.gov/air/criteria.html>.

EPA adopted a lower fine particulate NAAQS and revoked the annual PM₁₀ standard; both became effective December 17, 2006. They also lowered the ozone standard, effective May 27, 2008.

Air Quality Standards for Criteria Pollutants

Pollutant	Standard	Level
Ozone	The 3-year average of the 4 th highest daily maximum 8-hour average concentration cannot exceed the level measured at each monitor within an area over each year.	0.075 ppm
Particulate Matter (10 micrometers)	The 3-year annual average of the weighted annual mean concentration at each monitor within an area must not exceed the level	Revoked
	The 24-hour average cannot exceed the level more than once per year	154 µg/m ³
Particulate Matter (2.5 micrometers)	The 3-year annual average of the weighted annual mean concentrations cannot exceed the level	15.4 µg/m ³
	The 3-year average of the 98 th percentile (based on the number of samples taken) of the daily concentrations must not exceed the level	35.4 µg/m ³
Carbon Monoxide	The 1-hour average cannot exceed the level more than once per year	35.4 ppm
	The 8-hour average cannot exceed the level more than once per year	9.4 ppm
Sulfur Dioxide	Annual arithmetic mean of 1-hour averages cannot exceed the level	0.03 ppm
	24-hour average cannot exceed the level more than once per year	0.14 ppm
	The 3-hour average cannot exceed the level more than once per year	0.50 ppm
Lead	The quarterly average (by calendar) cannot exceed the level	1.5 µg/m ³
Nitrogen Dioxide	The annual mean of 1-hour averages cannot exceed the level	0.053 ppm

Note: Daily concentration is the 24-hour average, measured from midnight to midnight.



2007 Air Quality Data Summary

In some instances, comparison of numbers in this table with sources listed above may appear to be slightly off (for example, the carbon monoxide 8-hour standard 9.4 ppm listed in the table versus 9.0 ppm on the EPA Web site). These slight differences are due to a rounding convention adopted by EPA and the number of significant figures. The numbers shown on the table above are those used to determine if an area is in compliance, and are reflected in the graphs on the following pages.

The NAAQS for each pollutant may have different averaging times (for example, hourly and 8-hour averages). These different forms of the standard are created and enforced to address varied health impacts that happen as a result of a shorter, high-level exposure versus longer, low-level exposures. These differences are addressed pollutant-by-pollutant in the following sections, and additional information is on the EPA Web site. A distinction exists between “exceeding” and “violating” a standard; the two are not equivalent. This distinction is due to the nature of the standards. In most instances it is allowable for an area to exceed the standard a few times, to allow for possible unusual meteorological circumstances. For example, a carbon monoxide 8-hour average of 10 ppm clearly exceeds the standard; however, it does not violate the standard if it is the only exceedance that year (the standard allows for one exceedance).

The EPA standards typically apply to an ‘area’, which may be defined in different ways. Data are often presented for individual monitoring stations in the following sections because this provides more insight into regional differences in Idaho’s ambient air quality. The summaries that follow show how Idaho’s airsheds compared to the standards discussed above for the year 2007 and in many instances also incorporate the AQI and other measures of air quality where appropriate. The AQI color code shading is shown to aid interpretation of air quality, but does not imply whether or not standards were actually met for each pollutant. An airshed must satisfy the conditions in the above table to ensure compliance with the NAAQS.



2007 Air Quality Data Summary

Monitoring Network

The Idaho monitoring network is a composite of meteorological and pollutant-specific monitoring equipment. DEQ operates most of the monitors while several tribes operate monitors on tribal lands. Data from the network are either collected manually by field staff or sent directly to engineers and scientists through a telemetry network.

The table on the next page presents a summary of the monitoring stations used and parameters monitored during 2007. Some parameters were monitored for only part of the year.

The map on page 8 shows monitoring stations that were active in 2007. Monitoring stations are mainly located in high population areas; however, DEQ does monitor air quality in some rural areas. Some sites are selected to focus on the emissions of a single pollutant or group of sources (for example, near a high traffic volume or residential wood burning area). Monitor siting and monitoring objectives are discussed in the pollutant-specific sections of this report.

Criteria pollutants are measured using methods approved by EPA, to assess Idaho's compliance with National Ambient Air Quality Standards (NAAQS). In addition, some pollutants of particular interest are measured using more than one method. These additional methods help engineers and scientists to better understand the presence and behavior of these pollutants. The table on page seven lists the methods used for the various pollutants. It is noteworthy that the tapered element oscillating microbalance (TEOM) method (for particulate matter) is a continuous monitoring method. A TEOM measures mass concentrations at pre-set time intervals (e.g. hourly). The TEOM can also be accessed through telemetry for instantaneous PM concentrations. TEOMs enable real-time data interpretation and are discussed further in the particulate matter section of this report. Additional information on measurement methods is available at EPA's Web site: <http://www.epa.gov/ttn/amtic/>.

In addition to the criteria air pollutants described in this report, air toxics were monitored for two years at a Nampa site. Monitoring was terminated at this site on March 11, 2005. DEQ intends to resume air toxics monitoring some time in the future. For details on air toxics and chemical toxicity, visit the EPA web site at www.epa.gov/ttn/atw/index.html.

Particulate Monitoring

Coarse particulate (PM₁₀) and fine particulate (PM_{2.5}) are measured using a variety of methods in Idaho. EPA considers the federal reference method (FRM) to be the most accurate way to determine PM₁₀ and PM_{2.5} concentrations. This method involves pulling in air (at a given flow rate) and trapping particles of a certain size (PM₁₀ or PM_{2.5}) on a pre-weighed filter. The filter is then weighed again and the resulting mass is divided by volume (determined from flow rate and amount of time) to provide concentration. Particles on the filter can be later chemically analyzed and modeled for more information about the sources of particulate matter. Unfortunately, the FRM does not provide continuous or timely information. EPA considers the TEOM continuous method equivalent to the FRM (Federal Equivalent Method, or FEM) for PM₁₀, but not an equivalent method for PM_{2.5}.



2007 Air Quality Data Summary

DEQ uses the FRM as well as the TEOM continuous method to provide more time-resolved data (i.e. hourly averages).

Monitoring Network for 2007

Name	Location	PM ₁₀ FRM	PM ₁₀ TEOM	PM _{2.5} FRM	PM _{2.5} TEOM	O ₃	SO ₂	NO _x	CO	Met	PM _{2.5} CSM
Boise	Boise, various locations	●	●	●	●	●		●	●	●	
Coeur d'Alene	930 N 15 th , Lakes Middle School				●						
Franklin	Franklin, Idaho			●							
Grangeville	USFS Compound, East Grangeville				●					●	
Idaho Falls	North Holms & Pop Kroll, Idaho Falls				●						
Lancaster	Lancaster Road near Coeur d'Alene					●		●		●	
Lewiston	Sunset Park, Lewiston				●					●	
McCall	500 North Mission, McCall				●						
Meridian	St. Luke's				●	●		●			●
Moscow	1025 Plant Sciences Road				●					●	
Nampa	Nampa, various locations		●	●	●						
Parma	Parma				●						
P4/Monsanto	Soda Springs						●				
Pinehurst	106 Church Street, Pinehurst		●	●	●					●	
Pocatello	Pocatello, various locations	●	●	●	●		●			●	
Salmon	Salmon, various locations			●	●					●	
Sandpoint	Sandpoint		●		●					●	
St. Maries	St. Maries			●	●						
Twin Falls	Smith's Food Store, Twin Falls				●						
Wendell	Wendell, Gooding County									●	

Notes:

CO	Carbon Monoxide	PM _{2.5} FRM	Particulate Matter 2.5 micrometers (reference)
CSM	Chemical Speciation Monitor	PM _{2.5} TEOM	Particulate Matter 2.5 micrometers (TEOM continuous)
NO _x	Nitrogen Oxide	SO ₂	Sulfur Dioxide
O ₃	Ozone (May through September)	Tox	Urban Air Toxics
PM ₁₀ FRM	Particulate Matter 10 micrometers (reference)	●	Monitor Terminated in 2007
PM ₁₀ TEOM	Particulate Matter 10 micrometers (TEOM continuous)	●	NNU Precision Terminated 12/26/07
		●	Pinehurst Precision Started 12/2/2007



2007 Air Quality Data Summary


Monitoring Methods Used in 2007 in Idaho

Pollutant Code	Measurement	Method	Units
CO	Carbon Monoxide	Gas Nondispersive Infrared Radiation	Parts per Million
NO _x	Nitrogen Oxides (NO _x)	Chemiluminescence	Parts per Million
O ₃	Ozone	UV Absorption	Parts per Million
PM ₁₀ FRM	PM ₁₀ Reference	Reference - Hi Vol Andersen/ GMW 1200	Micrograms per Cubic Meter
PM ₁₀ TEOM	PM ₁₀ TEOM	R&P Mass Transducer	Micrograms per Cubic Meter
PM _{2.5} FRM	PM _{2.5} Reference	Reference—R&P Partisol 2025	Micrograms per Cubic Meter
PM _{2.5} TEOM	PM _{2.5} TEOM	R&P Mass Transducer	Micrograms per Cubic Meter
SO ₂	Sulfur Dioxide	UV Fluorescence	Parts per Million



2007 Air Quality Data Summary

IDAHO DEQ SFY 2007 Air Monitoring Network





Monitoring Results

Ozone

Ozone is typically a summertime air pollution problem which primarily forms when pollutants from cars and industrial sources (paints, solvents, gas vapors) react with sunlight. These pollutants are called ozone precursors and include VOCs and NO_x. Ozone can also be directly emitted by pollutant sources. Levels of ozone are usually highest in the afternoon because of the intense sunlight, warm temperatures, and the time required for ozone to form. These levels are highly affected by weather. DEQ monitored ozone from May through September in 2007, as this is the time period of concern for high ozone levels.

Ozone is considered beneficial in the upper atmosphere because it helps to protect the earth from the sun's rays; however, ozone formed at ground level is unhealthy. Elevated concentrations of ground-level ozone can cause reduced lung function, respiratory irritation, and can aggravate asthma. Ozone has also been linked to immune system effects (www.epa.gov/ttn/oarpg/naaqsfin/o3health.html). The damage ozone causes to the lungs typically heals within a few days, but repeated or prolonged exposure may cause permanent damage. People with respiratory conditions should limit outdoor exertion if ozone levels are high. Even healthy individuals may experience respiratory symptoms on a high-ozone day. Ground-level ozone can also damage agricultural crops and forests, interfering with their ability to photosynthesize and grow.

The monitoring stations measuring ozone are located in both urban and rural areas, although the precursor chemicals that react with sunlight to produce ozone are generated primarily in large metropolitan areas. Because summers in Idaho are normally hot and dry, ozone levels typically begin to rise in the late morning and then peak in the late afternoon and early evening. This phenomenon follows very closely with the time of day that the sun is the highest in the sky and temperatures are the hottest.

Graphs presented on pages 10-13 show trends in ozone levels in Idaho, reflecting both the AQI category and the NAAQS. The graphs on these pages present daily maximum eight-hour average data for the months of May through September. The shading on each graph corresponds to the AQI categories. It should be noted that AQI categories of orange and above indicate NAAQS exceedances.


The graph on page 14 shows monitoring data for each ozone monitoring station against the previous and the new eight-hour federal standard, and shows that the region has remained below the previous ozone standard since monitoring began. It also shows that the Treasure Valley is very close to the new standard. This means the three-year average of the fourth-highest eight-hour concentration will violate the new NAAQS if it exceeds 0.075 ppm (0.076 ppm or higher). The ozone standard is defined such that the three highest concentrations can exceed the level of the standard while still maintaining attainment. The averaging period from which compliance determinations to the new NAAQS has been defined by EPA to be 2006-2008. Values presented on the page 14 graph are three-year averages of the fourth-highest concentrations; the year on the x-axis represents the last year averaged. For example, concentrations shown for 2007 are an average of 2005, 2006, and 2007 concentrations.

For additional information on ozone, visit www.epa.gov/air/ozonepollution/. There is also additional information on ozone in question/answer format in the definitions section of this document.



2007 Air Quality Data Summary


Lancaster 8-Hour Ozone Daily Maximum 8-Hour Concentration





2007 Air Quality Data Summary

St. Luke's 8-Hour Ozone Daily Maximum 8-Hour Concentration




Note: St. Luke's ozone monitor experienced significant system failures during the monitoring season.



2007 Air Quality Data Summary

ITD 8-Hour Ozone Daily Maximum 8-Hour Concentration




Note: ITD ozone monitor experienced significant system failures during the monitoring season.



2007 Air Quality Data Summary

Whitney 8-Hour Ozone Daily Maximum 8-Hour Concentration






2007 Air Quality Data Summary

Idaho 8-Hour Ozone Averages

3-Year Average of 4th Highest Annual Concentration vs Federal Standard





2007 Air Quality Data Summary

Particulate Matter (10 micrometers)

Particulate matter (PM) includes both solid matter and liquid droplets suspended in the air. Particles smaller than 2.5 micrometers in diameter are called “fine” particles, or $PM_{2.5}$. Particles between 2.5 and 10 micrometers in diameter are called “coarse” particles. PM_{10} includes both fine and coarse particles. Coarse particles typically come from crushing or grinding operations and dust from roads. PM_{10} can aggravate respiratory conditions such as asthma. People with respiratory conditions should avoid outdoor exertion if PM_{10} levels are high.

The federal annual PM_{10} standard was revoked effective December 17, 2006 due to a lack of evidence linking health problems to long-term exposure to coarse particle pollution. The 24-hour standard was not changed. EPA may choose to replace the PM_{10} standard in the future with a $PM_{10-2.5}$ (PM_{coarse}) standard, ranging from diameters 2.5-10 micrometers. In 2007, Idaho had two areas that had previously exceeded federal PM_{10} standards. These areas are in Sandpoint and Pinehurst. Pocatello and Northern Ada County were formerly nonattainment areas but are now considered to be maintenance areas for PM_{10} .

Idaho monitors PM_{10} using both reference and continuous methods. The PM_{10} TEOM is a federal equivalent method. TEOM data is also used to determine compliance to the PM_{10} NAAQS. Reference and equivalent method results are shown in the following graphs. TEOM data is also used to determine the daily AQI and to inform the public of air quality values in near real-time via DEQ Web pages.



The graphs on the page 16 demonstrate that Idaho’s airsheds were in compliance for the daily NAAQS standards for PM_{10} in 2007. Pocatello G&G and Boise Mountain View School PM_{10} were measured using the federal reference method while Sandpoint, Pinehurst, the Boise Fire Station, and Nampa were measured with TEOMs, the federal equivalent method. The graph shows the rolling three (3) year average (year 2007 represents the average of 2005, 2006, 2007) of PM_{10} concentrations at individual monitoring stations. It is clear that all concentrations are below the NAAQS.

Maximum daily values (24-hour average) confirm that Idaho has generally shown a decrease since 1996, although the Nampa site is showing a three-year increase. Statistical summaries of reference and continuous method PM_{10} concentrations are provided in tables in the Appendix. The maximum PM_{10} measured in 2007 at the Nampa monitor exceeded the 24-hour NAAQS standard. However, the 24-hour PM_{10} NAAQS is only considered violated if there are more than three exceedances during the consecutive three (3) year period. For example, we could experience two (2) exceedances in year 1, none in year 2, and one (1) in year 3 and not violate the NAAQS.

For additional information on PM_{10} , visit www.epa.gov/oar/particlepollution/. More information on PM_{10} is also presented in question/answer format in the definitions section of this document.



2007 Air Quality Data Summary





2007 Air Quality Data Summary

Particulate Matter (2.5 micrometers)

Particles 2.5 micrometers in diameter or less are called “fine” particles, or $PM_{2.5}$. DEQ considers $PM_{2.5}$ to be one of the major air pollution concerns affecting a number of airsheds in Idaho. $PM_{2.5}$ generally comes from wood burning, agricultural burning and other area sources, as well as vehicle exhaust including cars, diesel trucks, and buses. Fine particulate can also be formed secondarily in the atmosphere by chemical reactions of pollutant gases.

Exposure to $PM_{2.5}$ can have serious health effects. Fine particles are most closely associated with increased respiratory disease, decreased lung function, and even premature death. Children, older adults, and people with some illnesses are more sensitive and more likely to develop heart or lung problems associated with $PM_{2.5}$. People with respiratory or heart disease, older adults, and children should avoid outdoor exertion if $PM_{2.5}$ levels are high. $PM_{2.5}$ also significantly affects visibility.

$PM_{2.5}$ is primarily measured using two different methods in Idaho, the federal reference method and the Tapered Element Oscillating Method (TEOM). The federal reference method is the method approved by EPA to determine $PM_{2.5}$ NAAQS compliance. This method involves pulling in air (at a given flow rate) and trapping particles of a certain size (in this case $PM_{2.5}$) on a pre-weighed filter. The filter is then used, weighed, and divided by volume (determined from flow rate and amount of time) to provide the concentration. Unfortunately, the reference method does not provide continuous or timely information. Thus, Idaho uses the TEOM method to provide more time-relevant data. The TEOM method uses measurement of mass to determine particulate matter present. A third method of $PM_{2.5}$ measurement is used during agricultural burning season; the nephelometer. These transportable instruments help DEQ estimate $PM_{2.5}$ concentrations during these activities.

The graphs in this section use data primarily from the federal reference method. The continuous data are from the TEOMs. The continuous methods are compared to the reference method values for a one year period and calculations are made to determine the degree of difference from the reference method. The differences are then applied to the current continuous values in an attempt to make them “reference method-like.” Data gathered by the TEOM or nephelometer method(s) cannot be used for NAAQS compliance determinations unless an agency demonstrates to EPA method equivalency by the recently promulgated “Approved Regional Method” process. DEQ has not begun this process.

The graphs on pages 19 and 20 show that all $PM_{2.5}$ monitors meet the annual NAAQS, and all but Pinehurst meet the daily NAAQS using the federal reference method. All of Idaho was designated attainment/unclassifiable for $PM_{2.5}$ in 2007. The graphs on pages 21 through 27 use data from the continuous samplers to display $PM_{2.5}$ with the AQI. Both types of graphs are discussed below.

The graph on page 19 shows the 2007 three-year average of the 98th percentile 24-hour (daily) averages at each monitoring station against the federal standard. The annual averages for 2001-2007 all fell well below the previous standard of $65 \mu\text{g}/\text{m}^3$. For 2007 the graph shows the three year average for Pinehurst violating the new NAAQS of $35 \mu\text{g}/\text{m}^3$. There are only two years of data for Franklin, but it was designated non-attainment because it is in the same airshed as Logan, Utah, which violated the 24-hr NAAQS.



2007 Air Quality Data Summary

The graph on page 20 shows the three-year average of the annual averages at each monitoring station, against the federal standard. It is easily seen that the annual standard of $15.4 \mu\text{g}/\text{m}^3$ was not exceeded at any of the monitoring stations.

Graphs on pages 21 through 26 show daily $\text{PM}_{2.5}$ concentrations measured at Idaho sites during 2007 using the TEOM continuous analyzers against a backdrop of AQI breakpoints. The highest concentration of $\text{PM}_{2.5}$ measured with the TEOMS in 2007 was $126.5 \mu\text{g}/\text{m}^3$, measured at Salmon during a natural wildland fire event. There were numerous summer fires near Salmon in 2007 and the Eastern Idaho graph reflects this. A few of the graphs show some blank periods with no concentrations. These are times when a TEOM was not functioning due to mechanical malfunctions.


For additional information on particulate matter, visit www.epa.gov/oar/particlepollution/. Information on $\text{PM}_{2.5}$ is also presented in a question/answer format in the definitions section of this document.



2007 Air Quality Data Summary

3-Yr Average 98% Daily PM_{2.5} 2001 - 2007

Federal Reference Method Monitors



Franklin and Salmon have been included for reference even though they do not yet have three years of data to average.

The Salmon value was during an exceptional event (unwanted wildfire) and DEQ has flagged it per EPA's exceptional events policy. If EPA agrees, this value will not be used for NAAQS compliance.


Monitors included for historical reference but are currently not operating include: Pocatello G&G, Sandpoint, Idaho Falls, Coeur d'Alene, Lewiston, and Twin Falls. Franklin and Salmon do not have three years of data, but are included for reference.



2007 Air Quality Data Summary

3-Year Average Annual Mean PM_{2.5} 2001 - 2007

Federal Reference Method Monitors




Franklin and Salmon have been included for reference even though they do not yet have three years of data to average.


Monitors included for historical reference but are currently not operating include: Pocatello G&G, Sandpoint, Idaho Falls, Coeur d'Alene, Lewiston, and Twin Falls. Franklin and Salmon do not have three years of data, but are included for reference.



2007 Air Quality Data Summary



Note that any gaps in TEOM continuous data are due to invalid data or system failure.






2007 Air Quality Data Summary

Coeur d'Alene


PM_{2.5} Daily Averages from Continuous Analyzers



Note that any gaps in TEOM continuous data are due to invalid data or system failure.


St. Maries

PM_{2.5} Daily Averages from Continuous Analyzers






2007 Air Quality Data Summary



Note that any gaps in TEOM continuous data are due to invalid data or system failure.






2007 Air Quality Data Summary

Lewiston


PM_{2.5} Daily Averages from Continuous Analyzers



Note that any gaps in TEOM continuous data are due to invalid data or system failure.


Moscow

PM_{2.5} Daily Averages from Continuous Analyzers






2007 Air Quality Data Summary




Note that any gaps in TEOM continuous data are due to invalid data or system failure.






2007 Air Quality Data Summary



Note that any gaps in TEOM continuous data are due to invalid data or system failure.





2007 Air Quality Data Summary

Carbon Monoxide

Carbon Monoxide (CO) is an odorless, colorless gas that can enter the bloodstream through the lungs and reduce the amount of oxygen that reaches organs and tissues. Carbon monoxide forms when the carbon in fuels doesn't burn completely. The majority of CO comes from vehicle exhaust. In cities, 85-95% of all CO emissions may come from motor vehicle exhaust.

Elevated levels of CO in the ambient air can occur in urban canyon areas with heavy traffic congestion. The highest levels of CO in the outside air typically occur during the colder months of the year when temperature inversions are more frequent. People with cardiovascular disease or respiratory problems might experience chest pain and increased cardiovascular symptoms, particularly while exercising, if CO levels are high. High levels of CO can affect alertness and vision even in healthy individuals.

CO monitoring stations are generally located in urban canyon areas with heavy traffic congestion. These include central business areas, roadsides, and shopping malls. Idaho currently only monitors CO in Boise as a condition of the CO Maintenance Plan.

The graph on page 28 shows the second highest eight-hour concentrations at Idaho's monitoring sites versus the NAAQS standard from 2001 through 2007. The second-highest concentration is displayed on these graphs because, under the federal rule, the eight-hour average can not be exceeded more than once per year (thus, choosing the second highest). These graphs confirm the general downward trend that CO is taking from the early 1990s to present. There were no eight-hour concentrations measured at any sites that exceeded the NAAQS standard of 9.4 ppm. The maximum eight-hour concentration for CO in 2007 was 1.7 ppm, well below the eight-hour standard. These data are provided in the Appendix.

The NAAQS also includes a 1-hour standard for CO of 35 ppm (can not be exceeded more than once a year). Measured 1-hour concentrations in Idaho are historically much lower than the 35 ppm standard, and therefore 1-hour CO trends were not graphed. The maximum and second-highest measured 1-hour CO in 2005 are 4.6 and 4.3 ppm, respectively. Additional 1-hour average CO data are provided in the Appendix.


For additional information on CO, visit www.epa.gov/air/urbanair/co/index.html. CO information is also provided in question/answer format in the definitions section of this document.



2007 Air Quality Data Summary

Carbon Monoxide (CO) for Idaho

2nd Highest 8-Hour Concentration vs Standard





2007 Air Quality Data Summary

Sulfur Dioxide

Sulfur dioxide (SO₂) is a colorless, reactive gas produced by burning fuels containing sulfur, such as coal and oil, and by industrial processes. Historically, the greatest sources of SO₂ were industrial facilities that derived their products from raw materials like metallic ore, coal, and crude oil, or that burned coal or oil to produce process heat (petroleum refineries, cement manufacturing, and metal processing facilities). Currently, on-road vehicles, marine craft, and diesel construction equipment also release significant SO₂ emissions to the air.

People with asthma who are active outdoors may experience bronchoconstriction, where symptoms include wheezing, shortness of breath, and tightening of the chest. People should limit outdoor exertion if SO₂ levels are high.

The graph on page 30 shows that Idaho is well below the annual standard for SO₂. The maximum measured SO₂ concentrations in 2007 were significantly below the federal standards as well. The graphs on pages 31 and 32 show the maximum 24-hour and 3-hour concentrations, respectively, at Idaho's monitoring sites. The maximum 24-hour and 3-hour averages were 0.090 ppm and 0.107 ppm, respectively. Note that the 2005-2007 Soda Springs monitor is at a different location than it was in 1999-2002. DEQ changed from population exposure monitoring to "hot spot" monitoring at Soda Springs. 'Hot spot' refers to monitoring that is designed to investigate pollution sources on a local scale. This allows for the assessment of air quality emanating from a point source, but rather than emissions being monitored directly from a stack or chimney, the air is measured as it moves towards areas where it may impact on human health or quality of life.


Additional SO₂ data are located in the Appendix, and information on SO₂ is available at www.epa.gov/air/urbanair/so2/index.html. SO₂ information is also provided in question/answer format in the definitions section of this document.



2007 Air Quality Data Summary

Sulfur Dioxide (SO₂)

Annual Average vs Standard






2007 Air Quality Data Summary

Sulfur Dioxide (SO₂)

Maximum 24-Hour Average vs Standard






2007 Air Quality Data Summary

Sulfur Dioxide (SO₂)

Maximum 3-Hour Average vs Standard





2007 Air Quality Data Summary

Lead

Lead is a highly toxic metal that was used for many years in household products, automobile fuel, and industrial chemicals. Airborne lead was associated primarily with automobile exhaust and lead smelters. The large reductions in lead emissions from motor vehicles have changed the nature of the air quality lead problem in the United States. Industrial processes, particularly primary and secondary lead smelters and battery manufacturers, are now responsible for most of the lead emissions.

People, animals, and fish are mainly exposed to lead by breathing and ingesting it in food, water, soil, or dust. Lead accumulates in the blood, bones, muscles, and fat. Infants and young children are especially sensitive to even low levels of lead. Lead can have health effects ranging from behavioral problems and learning disabilities to seizures and death.


According to EPA, the primary sources of lead exposure are lead-based paint, lead-contaminated dust, and lead-contaminated residual soils. Refer to the EPA Web site www.epa.gov/ttnatw01/hlthef/lead.html for ways to limit your exposure to these lead sources.

Lead has not been monitored in Idaho since 2002. With the phase-out of lead in fuel and the closure of the Bunker Hill lead smelter in Kellogg, airborne lead is no longer a public health concern in Idaho. The graph on page 34 is included to show the historical monitoring of airborne lead in Kellogg.

For additional information on lead, visit www.epa.gov/air/lead/. Lead information is also available in a question/answer format in the definitions section of this document.



2007 Air Quality Data Summary





Nitrogen Dioxide

Nitrogen dioxide (NO₂) is a reddish brown, highly reactive gas that forms from the reaction of nitrogen oxide (NO) and oxygen in the atmosphere. The term "NO_x", which is frequently used, refers to both NO and NO₂. NO₂ will react with VOCs and can result in the formation of ozone. On-road vehicles like trucks and automobiles are the major sources of NO_x. Industrial boilers and processes, home heaters, and gas stoves can also produce NO_x. NO₂ pollution is greatest in cold weather.

NO₂ can cause respiratory symptoms such as coughing, wheezing, and shortness of breath in people with respiratory diseases such as asthma. Long-term exposure can lead to respiratory infections.

Motor vehicle manufacturers have been required to reduce NO_x emissions from cars and trucks since the 1970s. NO_x is not considered a significant pollution problem in Idaho. In 2007, DEQ maintained three monitoring sites for nitrogen dioxide: Lancaster near Coeur d'Alene; Meridian; and Boise. The monitoring objective is primarily to assess ambient NO_x concentrations for evaluating ozone formation processes during the ozone season.


The maximum 1-hour average of NO₂ measured in 2007 was 0.033 ppm. The averages observed have consistently been well below the NAAQS annual standard, as shown in the graph on page 36 and in data in the Appendix. However, these averages cannot be used to assess NAAQS compliance since the monitors were not operated for the entire year.

For additional information on NO₂, visit www.epa.gov/air/urbanair/nox/index.html.



2007 Air Quality Data Summary

Idaho Nitrogen Dioxide (NO₂) Annual 1-Hour Average vs Standard





Air Quality Index

The AQI is reported according to a 500-point scale for each of the major criteria air pollutants: ozone, particulate matter (PM_{2.5} & PM₁₀), carbon monoxide, nitrogen dioxide, and sulfur dioxide. The “worst denominator” determines the ranking. For example, if an area has a carbon monoxide value of 132 on a given day and all other pollutants are below 50, the AQI for that day would be 132. The AQI scale breaks down into six categories. Each category has a corresponding color, shown below, with [pollution concentration breakpoints](#) for each category.

Levels of Health Concern	Numeric Value	Meaning
Good	0-50	Air quality is considered satisfactory, and air pollution poses little or no risk.
Moderate	51-100	Air quality is acceptable; however, for some pollutants there may be a moderate health concern for a very small number of people who are unusually sensitive to air pollution.
Unhealthy for Sensitive Groups	101-150	Members of sensitive groups may experience health effects. The general public is not likely to be affected.
Unhealthy	151-200	Everyone may begin to experience health effects; members of sensitive groups may experience more serious health effects.
Very Unhealthy	201-300	Health alert: everyone may experience more serious health effects.
Hazardous	300-500	Health warnings of emergency conditions. The entire population is more likely to be affected.


The AQI is a national index, so the values and colors used to show local air quality and the associated level of health concern will be the same everywhere in the U.S.A. The number of “good” air quality days continues to dominate regionally in Idaho. However, there were brief periods when the air quality degraded into “moderate,” “unhealthy for sensitive groups,” “unhealthy,” and “very unhealthy.” The table presented on page 48 shows the AQI breakdown by percentage in each category for the year.

The graph on page 38 presents the annual number of “good” days for several Idaho counties. The number of “good” days has remained relatively high over the last few years for each county. Current reduction in the number of “good” days cannot be directly compared with the numbers before 1999. In that year, PM_{2.5} was added to the index and the “unhealthy” category was divided into “unhealthy” and “unhealthy for sensitive groups.” In 2002 ozone monitoring was added to the AQI calculation in the Treasure Valley with ozone contributing to the number of moderate days in the following years. Ozone monitoring began at the Lancaster site near Coeur d’Alene in 2005.

The AQI Graphs on pages 39-47 present the distribution of AQI categories recorded for nine Idaho counties. The AQI data summaries for each county that support the graphs are located in the Appendix.




2007 Air Quality Data Summary





2007 Air Quality Data Summary


Air Quality for Ada County





2007 Air Quality Data Summary


Air Quality for Bannock County





2007 Air Quality Data Summary


Air Quality for Bonneville County





2007 Air Quality Data Summary


Air Quality for Canyon County





2007 Air Quality Data Summary


Air Quality for Franklin County





2007 Air Quality Data Summary


Air Quality for Kootenai County





2007 Air Quality Data Summary


Air Quality for Nez Perce County





2007 Air Quality Data Summary


Air Quality for Shoshone County





2007 Air Quality Data Summary

Air Quality for Twin Falls County





2007 Air Quality Data Summary

The number of “good” AQI days continues to dominate regionally in Idaho; however, there were brief periods when the air quality degraded into one of the other categories. The table below shows the AQI breakdown by percentage in each category where air quality is monitored. In 2007, the highest AQI value of 223 was recorded in Lemhi County for PM_{2.5}. This value was in the very unhealthy range.

2007		AQI Rating # Days				
County	# AQI Days	Good	Moderate	Unhealthy for Sensitive Groups	Unhealthy	Highest AQI
Ada	365	251	102	11	1	163
Bannock	359	353	6			90
Benewah	303	277	26			96
Bonner	305	305				49
Bonneville	276	264	12			73
Butte	222	207	15			87
Canyon	362	289	71	2		123
Caribou	347	346	1			76
Franklin	211	199	11	1		104
Idaho	340	322	17		1	163
Kootenai	357	303	54			90
Latah	353	347	6			73
Lemhi	316	272	35	6	3	223
Nez Perce	357	348	9			67
Power	339	313	26			81
Shoshone	361	237	114	10		133
Twin Falls	295	275	20			71
Valley	167	120	40	6	1	153

There were no NAAQS violations in 2007 for any pollutants. In most cases, pollutant concentrations fell well below standards.



Impaired Air Quality

Winter Burn Bans

Idaho has a winter-impaired air quality program primarily targeting sources of particulate matter from wood stoves and fireplaces. Idaho's program is generally implemented through local ordinances in those areas that have historically had winter inversion problems. Generally, these ordinances specify that a wood burning ban will be declared whenever DEQ reports an AQI value of 74 or greater for any pollutant measured by an approved monitor and air stagnation conditions are forecasted to continue for at least 24 hours. In some areas, open burning (even if a valid permit has been issued) is prohibited when DEQ reports an AQI value of 60 or higher for any pollutant measured in the city/region/airshed.

The DEQ online [Current Air Quality Report](#) lists the daily air quality in many cities and regions in Idaho. Each report will list the pollutant being monitored, the AQI, and burn restrictions, if any, for the day. Anyone wanting to know if they can burn can go to this site to see what the forecast is for their area.

DEQ will issue an air quality advisory for specific locations between 2:00 and 4:00 p.m. on days where air quality is forecast to be poor. The advisories will be for the expected conditions the next day. Updates will be sent out on weekend days at the same times if conditions are expected to be poor. These advisories will be provided to local media outlets and to others through email notification lists.

Summer Ozone Alerts

DEQ forecasts pollution conditions for ozone in the Treasure Valley using pollutant monitoring data and meteorological information. Because ozone needs heat and sunlight to form, it is considered a summertime problem and is only monitored from May 1 through September 30. Ozone pollution can rise to very high levels when the valley experiences hot days with few clouds in the sky. The Treasure Valley tends to see daily ozone levels that begin to rise in the late morning and peak in the late afternoon and early evening. This phenomenon follows very closely with the time of day that the sun is the highest in the sky through the time temperatures are the hottest. Since we have no control over our weather characteristics, we have to focus on controlling what we put into our air. Under yellow or moderate alerts, the public is requested to change certain behaviors to prevent further deterioration of air quality. These alerts will be reported to local media outlets and to others through an email notification list.



DEFINITIONS

General Definitions

Air Toxics

Air toxics are broadly defined as almost 700 pollutants that DEQ considers to be potentially harmful to human health and the environment. These pollutants are listed in the Idaho air rules in IDAPA 58.01.01.585 and 586 (<http://adm.idaho.gov/adminrules/rules/idapa58/0101.pdf>). Hazardous air pollutants (see below) are included in this list to identify them as a subset of air toxics.

Criteria Air Pollutant (CAP)

The Clean Air Act of 1970 defined six *criteria pollutants* and established ambient concentrations of each to protect public health. EPA periodically has revised the original concentration limits and methods of measurement, most recently in 2008. [See page 3 for the list and the allowed ambient concentrations.](#)

Hazardous Air Pollutant (HAP)

A *hazardous air pollutant* is an air contaminant identified as toxic in the Federal Clean Air Act, Section 112(b). 188 pollutants are currently listed as HAPs. They are listed by EPA at <http://www.epa.gov/ttn/atw/188polls.html>.

Temperature Inversions

The earth gains and loses most of its energy at its surface. It is warmed by solar heating during the day and cooled by radiation emissions at night. During the late morning and afternoon hours, the air near the surface is warmer than the air aloft and allows for good pollutant dispersion (vertical mixing may be 1,500 meters or more). At night with clear skies, the surface radiates heat into outer space, creating cooler air at the surface and warmer air aloft. Warmer air above cooler air (temperature inversion) is a stable condition and limits the upward movement of pollution because the warmer air acts as a barrier. With little or no wind, pollutants are trapped near the surface (vertical mixing may be 200 meters or less) and can reach high levels of concentration.

Volatile Organic Compound (VOC)

An organic compound that participates in atmospheric photochemical reactions. This excludes all compounds determined to have negligible photochemical reactivity by EPA and listed in 40 CFR 51.100(s) in effect July 1, 1998.

Visibility/Regional Haze

Visibility is often explained in terms of visual range and light extinction. *Visual range* is the maximum distance—usually miles or kilometers—that you can see a black object against the horizon. *Light extinction* is the sum of light scattering and light absorption by fine particles and gases in the atmosphere. The more light extinction you have, the shorter your visual range will be. Reduced visibility (or visual range) is caused by weather (clouds, fog, and rain) and air pollution (fine particles and gases). The major pollution contributor to reduced visibility is fine particulate matter (PM_{2.5}) emissions, which are transported aloft and may remain suspended for a week or longer. Fine particles have a greater



2007 Air Quality Data Summary

impact than coarse particles at locations far from the emitting source because they remain suspended in the atmosphere longer and travel farther. $PM_{2.5}$ also presents some of the most serious health hazards to the public, so you can roughly assume that the worse the visibility, the unhealthier the air is to breathe.

Pollution Sources

Area Sources

Categories of pollution sources, in which each individual industrial source emits pollutants below the thresholds for a point source facility designation, and other categories that are a result of human activities and are best estimated at a county level in association with population numbers. For example, natural gas use for home heating, gas stoves, or woodstoves.

Biogenics

Natural sources such as trees, plants, grass, crops, and soils. The worldwide emissions rate of these natural hydrocarbons has been estimated to exceed that of non-methane hydrocarbons originating from human sources. Isoprene, one of the major constituents of biogenic emissions, is very photoreactive and makes biogenic VOCs a contributor in the formation of ozone.

Emission Factor

A value derived from source tests, material balance calculations, or engineering comparisons with similar processes. Used to estimate emissions from process quantities.

Non-road Mobile Sources

Farm vehicles, on-site construction/industrial vehicles, logging equipment, small marine craft, aircraft, trains, lawn and garden equipment, and off-road trail machines.

On-road Mobile Sources

Cars, trucks, sport utility vehicles, motorcycles and buses.

Point Sources

For the every-third-year statewide emissions inventory, point sources are defined as facilities that have actual annual air pollutant emissions equal to or exceeding 1000 tons per year of CO; 100 tons per year of NO_x , PM_{10} , $PM_{2.5}$, SO_x , or VOCs; or 5 tons per year of lead.

Registered Facility

The total of all pollutant-emitting activities located on adjacent or contiguous properties owned or operated by one person or corporate entity. It includes all of the pollutant-emitting buildings, processes, structures, equipment, control apparatuses, and storage areas at a facility.



Criteria Air Pollutants

Ozone (O₃)

- **What is it?**

Ozone, a bluish-colored gas molecule with a strong odor, is composed of three atoms of oxygen. In the upper atmosphere ozone occurs naturally and partially absorbs the sun's harmful ultraviolet rays. Ozone at ground level is a summertime air pollution problem.

- **How is it caused?**

Ozone forms when photochemical pollutants from cars, trucks, and industrial sources react with sunlight. Ozone-forming pollutants include NO_x and VOCs; even gasoline-powered yard equipment, paints, solvents, and off-road vehicle motors contribute.

- **When does it happen?**

Ozone pollution is most common in the summer months, when sunlight and stable atmospheric conditions occur. Ozone levels are usually highest in the afternoon, as sunlight photochemically transforms NO_x and VOCs into ozone.

- **Who is affected?**

Adults and children who are active outdoors, people with respiratory disease such as asthma, and people with unusual sensitivity to ozone. During physical activity, ozone penetrates deeper into the lungs and can do more damage.

Ozone is a very reactive gas. For this reason, high concentrations of ozone can cause respiratory distress and disease in humans, decreased yields of agricultural crops and forests, and damage to some rubber products, plastics, and paints used outdoors. National crop losses from ozone exposure are estimated at \$3 billion to \$5 billion annually. Forest losses are harder to estimate.

- **What are the health effects?**

Ozone can cause coughing and throat irritation, make deep vigorous breathing more difficult, and increase the chance of respiratory infections. It increases sensitivity to allergens and can trigger asthma attacks. The damage it causes to the lungs heals within a few days, but repeated or prolonged exposure may cause permanent damage.

- **What can I do about it?**

If ozone levels are high and you have a respiratory condition or are normally active outdoors, try to limit your outdoor exertion.

In the United States, management of ozone and other photochemical oxidants has been a major goal of federal and state clean air legislation (Clean Air Act). Although many of the pollution control efforts required by the CAA have been implemented, efforts to decrease ozone pollution have been only partially successful.

- **Where is it measured?**

Unlike other pollutants monitored here in Idaho, ozone is formed when precursor compounds react in the atmosphere. Winds transport ozone and precursor emissions from one area to another. For the Treasure Valley, ozone precursors are emitted into the air in urban areas of the airshed and



2007 Air Quality Data Summary

subsequently travel southeasterly to more rural areas as they react to form ozone. As a result, for the Treasure Valley airshed, DEQ has monitors in various locations. Another ozone monitor has been running in the Coeur d'Alene area since 2005.

Particulate Matter (PM_{2.5} and PM₁₀)

- **What is it?**

Particulate matter (PM) includes both solid matter and liquid droplets suspended in the air. Particles smaller than 2.5 micrometers in diameter are called “fine” particles, or PM_{2.5}. Particles between 2.5 and 10 micrometers in diameter are called “coarse” particles. PM₁₀ includes both fine and coarse particles. DEQ considers PM_{2.5} to be one of the major air pollution concerns affecting our state.

- **How is it caused?**

PM_{2.5} comes from all types of combustion, including cars, diesel trucks, power plants, wood burning, and from some industrial processes. It can also be formed in the atmosphere by chemical reactions of pollutant gases. The “coarse” particles in PM₁₀ typically come from crushing or grinding operations and dust from roads.

- **When does it happen?**

Daily PM_{2.5} trends in urbanized areas suggest that PM_{2.5} levels peak in association with traffic flow and rush hour periods. Periods of stagnate weather patterns, such as when surface inversions typically occur, contribute to elevated PM_{2.5} trends.

- **Who is affected?**

People with asthma and heart or lung disease, the elderly, and children. PM_{2.5} also significantly affects visibility.

- **What are the health effects?**

Fine particulates (PM_{2.5}) pose a greater risk to human health than coarse particulates, because they penetrate deeper into the respiratory system. PM_{2.5} exposure can have serious health effects. People with heart or lung diseases are at increased risk of attacks or premature death. Children and the elderly are more likely to develop heart or lung problems. PM₁₀ can aggravate respiratory conditions such as asthma.

- **What can I do about it?**

If PM_{2.5} levels are high, people with respiratory or heart disease, the elderly, and children should avoid outdoor exertion. If PM₁₀ levels are high, people with respiratory conditions should avoid outdoor exertion.

- **Where is it measured?**

Due to the health risks associated with PM, both PM_{2.5} and PM₁₀ are monitored in various population-oriented locations throughout Idaho.

Carbon Monoxide (CO)

- **What is it?**

CO is an odorless, colorless gas that can enter the bloodstream through the lungs and reduce the amount of oxygen that reaches organs and tissues.



2007 Air Quality Data Summary

- **How is it caused?**
Carbon monoxide forms when the carbon in fuels doesn't burn completely. Vehicle exhaust contributes 60% of all CO. In cities, that can be a 95% contribution.
- **When does it happen?**
CO pollution is at its worst in cold weather because fuels burn less efficiently in low temperatures. CO levels usually peak during morning and evening rush hours.
- **Who is affected?**
People with cardiovascular disease, such as angina, or cardiovascular or respiratory problems; also fetuses and young infants.
- **What are the health effects?**
Chest pain and increased cardiovascular symptoms, particularly while exercising. High levels of CO can even affect alertness and vision in healthy individuals.
- **What can I do about it?**
If CO levels are high, limit exertion and avoid sources of CO such as heavy traffic.
- **Where is it measured?**
CO monitoring stations are located in urban canyon areas with heavy traffic congestion. These include central business areas, roadsides, and shopping malls.

Sulfur Dioxide (SO₂)

- **What is it?**
Sulfur dioxide is a colorless, reactive gas.
- **How is it caused?**
SO₂ is produced by burning sulfur-containing fuels such as coal and oil and by some industrial processes.
- **Where does it happen?**
The highest concentrations of SO₂ are usually near large industrial sources.
- **Who is affected?**
People with asthma who are active outdoors.
- **What are the health effects?**
Bronchoconstriction, which can cause wheezing, shortness of breath, and tightening of the chest. When exposure to SO₂ ends, the symptoms should clear up within an hour.
- **What can I do about it?**
If SO₂ levels are high, limit your outdoor exertion.
- **Where is it measured?**
Because the large primary sources of SO₂ in Idaho are industrial, DEQ monitors for SO₂ near large facilities with high SO₂ emissions. The only monitors running in 2007 were in Pocatello and Soda Springs.

Lead (Pb)



2007 Air Quality Data Summary

- **What is it?**
Lead is a highly toxic metal that was used for many years in household products, automobile fuel, and industrial chemicals.
- **How is it caused?**
Locally, airborne lead is associated primarily with automobile exhaust and lead smelters. Since the phase-out of lead in fuels, cars and trucks are no longer a significant source of lead. Also, the Kellogg Bunker Hill Mine ceased operations in 1981.
- **When does it happen?**
Lead concentrations are likely to be highest near sources where current or former lead smelting/processing operations caused particle fallout, especially in nearby soils such as unpaved parking lots.
- **Who is affected?**
Everyone. Children six years and younger are most at risk.
- **What are the health effects?**
Lead can have health effects ranging from behavioral problems and learning disabilities to seizures and death.
- **What can I do about it?**
According to EPA, the primary sources of lead exposure are lead-based paint, lead-contaminated dust, and lead-contaminated residual soils. Refer to EPA's Web site at <http://www.epa.gov/ttn/atw/hlthef/lead.html> for ways to limit your exposure to these lead sources.
- **Where is it measured?**
Due to the phase-out of leaded fuels and the closure of Idaho's only lead smelter in 1981, DEQ no longer monitors for airborne lead. Historical monitoring was continued until 2002 but was discontinued due to the low levels being measured.

Nitrogen Dioxide (NO₂)

- **What is it?**
Nitrogen dioxide (NO₂) is a reddish brown, highly reactive gas that forms from the reaction of nitrogen oxide (NO) and oxygen in the atmosphere. NO₂ will react with VOCs and can result in the formation of ozone.
- **How is it caused?**
High temperature combustion sources such as power plants and automobiles are major producers of NO. Home heaters and gas stoves can also produce NO.
- **When does it happen?**
NO₂ pollution is greatest in cold weather. It follows a similar trend to CO.
- **Who is affected?**
People with respiratory diseases such as asthma; also children.



2007 Air Quality Data Summary

- **What are the health effects?**

NO₂ can cause respiratory symptoms such as coughing, wheezing, and shortness of breath. Long-term exposure can lead to respiratory infections.

- **What can I do about it?**

Since the 1970s, motor vehicle manufacturers have been required to reduce NO emissions from cars and trucks. It is not a significant pollution problem in Idaho.

- **Where is it measured?**

NO₂ is not a major concern in Idaho. It was measured during 2007 at the Lancaster site near Coeur d'Alene, Boise, and Meridian. It is now only monitored during the ozone season.

1410 N. Hilton
Boise, ID 83706

DEPARTMENT OF ENVIRONMENTAL QUALITY

www.deq.idaho.gov

Appendix



2007 Air Quality Data Summary Appendix

Calculation and Breakpoints for the Air Quality Index (AQI)

Breakpoints for Criteria Pollutants							AQI Categories	
O ₃ (ppm) 8-hour	O ₃ (ppm) 1-hour ^a	PM _{2.5} (µg/m ³)	PM ₁₀ (µg/m ³)	CO (ppm)	SO ₂ (ppm)	NO ₂ (ppm)	AQI value	Category
0.000–0.059	—	0.0–15.4	0–54	0.0–4.4	0.000–0.034	(b)	0–50	Good
0.060–0.075	—	15.5–35.4	55–154	4.5–9.4	0.035–0.144	(b)	51–100	Moderate
0.076–0.095	0.125–0.164	35.5–65.4	155–254	9.5–12.4	0.145–0.224	(b)	101–150	Unhealthy for sensitive groups
0.096–0.115	0.165–0.204	65.5–150.4	255–354	12.5–15.4	0.225–0.304	(b)	151–200	Unhealthy
0.116–0.374	0.205–0.404	150.5–250.4	355–424	15.5–30.4	0.305–0.604	0.65–1.24	201–300	Very unhealthy
(c)	0.405–0.504	250.5–350.4	425–504	30.5–40.4	0.605–0.804	1.25–1.64	301–400	Hazardous
(c)	0.505–0.604	350.4–500.4	505–604	40.5–50.4	0.805–1.004	1.65–2.04	401–500	

- a Areas are generally required to report the AQI based on 8-hour ozone values. However, there are a small number of areas where an AQI based on 1-hour ozone values would be safer. In these cases, in addition to calculating the 8-hour ozone value, the 1-hour ozone value may be calculated, and the greater of the two values reported.
- b NO₂ has no short-term National Ambient Air Quality Standard (NAAQS) and can generate an AQI only above a value of 200.
- c 8-hour O₃ values do not define higher AQI values (above 300). AQI values above 300 are calculated with 1-hour O₃ concentrations.

For more detailed information about the AQI and the pollutants it measures, go to www.epa.gov/airnow/aqibroch



2007 Air Quality Data Summary Appendix

- * US EPA - AirData Air Quality Index Report
- * Tuesday 15-Jul-2008 at 5:12:1 PM (USA Eastern time zone)
- * Geographic Area: Idaho
- * Pollutant: See <http://www.epa.gov/airnow/aqibroch/>
- * Year: 2007

Data Year	County	# AQI data days	Number of Days AQI was:				Max AQI	90th % AQI	Median AQI	Number of Days Main AQI Pollutant was:					
			Good	Mod	Unhealthy Sensitive Groups	Un-healthy				CO	NO2	O3	SO2	PM2.5	PM10
2007	Ada	365	251	102	11	1	163	74	41	13		128		185	39
2007	Bannock	359	353	6			90	34	17				148	102	109
2007	Benewah	303	277	26			96	44	21					303	
2007	Bonner	305	305				49	25	13						305
2007	Bonneville	276	264	12			73	41	20					276	
2007	Butte	222	207	15			87	50	38			222			
2007	Canyon	362	289	71	2		123	59	31					284	78
2007	Caribou	347	346	1			76	3	1				347		
2007	Franklin	211	199	11	1		104	39	16					211	
2007	Idaho	340	322	17		1	163	34	12					340	
2007	Kootenai	357	303	54			90	55	35			146		211	
2007	Latah	353	347	6			73	30	11					353	
2007	Lemhi	316	272	35	6	3	223	58	20					316	
2007	Nez Perce	357	348	9			67	38	15					340	17
2007	Power	339	313	26			81	48	25					194	145
2007	Shoshone	361	237	114	10		133	75	36					345	16
2007	Twin Falls	295	275	20			71	45	24					295	
2007	Valley	167	120	40	6	1	153	72	35					167	



2007 Air Quality Data Summary Appendix

- * US EPA - AirData Monitor Values Report - Criteria Air Pollutants
- * Wednesday 16-Jul-2008 at 11:18:10 AM (USA Eastern time zone)
- * Geographic Area: Idaho
- * Pollutant: Particulate (size < 2.5 micrometers)
- * Year: 2007

Data Year	County	24-Hour PM2.5							Annual Mean	Annual # Exceed	Monitor Number	Site ID	Site Address	City
		# Obs	1st Max	2nd Max	3rd Max	4th Max	98th Pct	# Ex-ceed						
2007	Ada	42	21	17	17	17	21	0	6.8	0	1	160010010	520 S. Eagle Road, Meridian	Meridian
2007	Ada	89	44	28	24	22	28	0	7.9	0	1	160010011	Mtn View School/3500 Carbarton Lane	Boise
2007	Benewah	113	39	31	29	26	29	0	9.9	0	1	160090010	9th And Center	St. Maries
2007	Benewah	27	14	9	9	8	14	0	5.9	0	1	160090011	850 A Street, Plummer	Plummer
2007	Canyon	19	28	17	15	12	28	0	9.3	0	2	160270004	Northwest Nazarine College (Nnc)	Nampa
2007	Canyon	243	52	29	27	26	26	0	7.6	0	1	160270004	Northwest Nazarine College (Nnc)	Nampa
2007	Franklin	211	42	36	33	29	27	0	6.9	0	1	160410001	Franklin - Water Treatment Facility At E	Franklin
2007	Idaho	35	36	25	19	19	36	1	10	0	2	160490003	Intersection Of Apple And Pine, Kamiah	
2007	Idaho	58	38	36	26	25	36	1	9.4	0	1	160490003	Intersection Of Apple And Pine, Kamiah	
2007	Lemhi	84	172	66	60	48	66	1	12.1	0	1	160590004	N. Charles St.	Salmon
2007	Shoshone	113	41	32	32	32	32	0	11.8	0	1	160790017	Pinehurst/Pinehurst School, Pinehurst	Pinehurst



2007 Air Quality Data Summary Appendix

- * US EPA - AirData Monitor Values Report - Criteria Air Pollutants
- * Wednesday 16-Jul-2008 at 1:10:52 PM (USA Eastern time zone)
- * Geographic Area: Idaho
- * Pollutant: Particulate (size < 10 micrometers)
- * Year: 2007

Data Year	County	24-Hour PM10						Annual Mean		Mon #	Site ID	Site Address	City
		# Obs	1st Max	2nd Max	3rd Max	4th Max	# Exceed-Actual	Annual Mean	# Exceed				
2007	Ada	343	88	79	77	69	0	24	0	3	160010009	Fire Station #5/16th & Front	Boise
2007	Ada	51	62	47	44	41	0	23	0	1	160010011	Mtn View School/3500 Carbarton Lane	Boise
2007	Bannock	91	46	31	28	27	0	11	0	3	160050015	G&G/Corner Of Garret & Gould	Pocatello
2007	Bannock	58	55	48	44	40	0	23	0	2	160050015	G&G/Corner Of Garret & Gould	Pocatello
2007	Bannock	81	51	51	45	44	0	22	0	1	160050015	G&G/Corner Of Garret & Gould	Pocatello
2007	Bonner	305	53	48	41	41	0	16	0	1	160170004	310 South Division Street	Sandpoint
2007	Canyon	329	175	82	79	71	1	25	0	2	160270002	Nampa Fire Stn/923 1st St	Nampa
2007	Nez Perce	20	47	38	37	33	0	27	0	2	160690006	Isp Bldg/2700 N/S Hwy 12-95	Lewiston
2007	Nez Perce	12	35	34	34	29	0	22	0	3	160690009	State Office Bldg/1118 F St	Lewiston
2007	Nez Perce	15	39	34	22	21	0	20	0	1	160690012	1200 29th Street	Lewiston
2007	Nez Perce	20	33	23	22	21	0	17	0	1	160690013	260 Bever Grade, Lapwai	
2007	Nez Perce	15	22	20	18	16	0	13	0	2	160690222	17500 Nez Perce Road	
2007	Nez Perce	19	45	29	23	22	0	16	0	1	160690222	17500 Nez Perce Road	
2007	Power	327	82	76	70	70	0	22	0	3	160770011	S Of Hwy 30 And E Of Weaver Rd	
2007	Power	59	57	52	39	36	0	19	0	2	160770011	S Of Hwy 30 And E Of Weaver Rd	
2007	Power	114	66	49	44	43	0	20	0	1	160770011	S Of Hwy 30 And E Of Weaver Rd	
2007	Shoshone	344	85	49	46	44	0	18	0	3	160790017	Pinehurst/Pinehurst School, Pinehurst	Pinehurst



2007 Air Quality Data Summary Appendix

- * US EPA - AirData Monitor Values Report - Criteria Air Pollutants
- * Wednesday 16-Jul-2008 at 10:31:32 AM (USA Eastern time zone)
- * Geographic Area: Idaho
- * Pollutant: Ozone
- * Year: 2007

Data Year	County	8-Hour Ozone								Monitor Number	Site ID	Site Address	City
		1st Max	2nd Max	3rd Max	4th Max	Days >Std	Required Days	# Days	% Days				
2007	Ada	0.074	0.069	0.069	0.068	0	153	59	39	1	160010010	520 S. Eagle Road, Meridian	Meridian
2007	Ada	0.082	0.081	0.080	0.080	6	153	89	58	1	160010019	3311 W. State Street, Boise	Boise
2007	Ada	0.086	0.085	0.078	0.078	5	153	128	84	1	160010030	Whitney Elementary School	Boise
2007	Butte	0.071	0.068	0.067	0.067	0	153	129	84	1	160230101	Craters Of The Moon National Mon, Idaho	COMNM
2007	Kootenai	0.072	0.071	0.071	0.067	0	153	146	95	1	160550003	North Of Lancaster Road - Near Hayden, I	Lancaster



2007 Air Quality Data Summary Appendix

- * US EPA - AirData Monitor Values Report - Criteria Air Pollutants
- * Tuesday 15-Jul-2008 at 5:35:18 PM (USA Eastern time zone)
- * Geographic Area: Idaho
- * Pollutant: Carbon Monoxide
- * Year: 2007

Data Year	County	1-Hour CO			8-Hour CO			Monitor #	Site ID	Site Address	City	
		# Obs	1st Max	2nd Max	# Exceed	1st Max	2nd Max					# Exceed
2007	Ada Co	6231	4.6	4.3	0	1.7	1.6	0	1	160010014	Eastman Bldg/166 N. 9th St	Boise



2007 Air Quality Data Summary Appendix

- * US EPA - AirData Monitor Values Report - Criteria Air Pollutants
- * Wednesday 16-Jul-2008 at 11:3:29 AM (USA Eastern time zone)
- * Geographic Area: Idaho
- * Pollutant: Nitrogen Dioxide
- * Year: 2007

Data Year	County	1-Hour NO2			Annual NO2		Monitor #	Site ID	Site Address	City
		# Obs	1st Max	2nd Max	Mean	# Exceed				
2007	Ada	1170	0.003	0.003	0.003	0	1	160010010	520 S. Eagle Road, Meridian	Meridian
2007	Ada	1466	0.030	0.025	0.004	0	1	160010019	3311 W. State Street, Boise	Boise
2007	Kootenai	3826	0.033	0.032	0.006	0	1	160550003	North Of Lancaster Road - Near Hayden, I	Hayden



2007 Air Quality Data Summary Appendix

- * US EPA - AirData Monitor Values Report - Criteria Air Pollutants
- * Wednesday 16-Jul-2008 at 11:10:5 AM (USA Eastern time zone)
- * Geographic Area: Idaho
- * Pollutant: Sulfur Dioxide
- * Year: 2007

Data Year	County	1-Hr SO2			3-Hr SO2			24-Hr SO2			Annual SO2		Mon #	Site ID	Site Address
		# Obs	1st Max	2nd Max	1st Max	2nd Max	# Exce	1st Max	2nd Max	# Excee	Mean	# Exce			
2007	Bannock	8431	0.085	0.074	0.052	0.051	0	0.026	0.024	0	0.006	0	2	160050004	Stp/Batiste & Chubbuck Rd
2007	Caribou	8342	0.121	0.105	0.107	0.09	0	0.090	0.005	0	0.002	0	1	160290031	5 Mile Road



2007 Air Quality Data Summary Appendix

US EPA - AirData Monitor Values Report - Criteria Air Pollutants
Friday, 21-Oct-2005 at 11:25:31 AM (USA Eastern time zone)
Geographic Area: Idaho
Pollutant: Lead
Year: 1999 - 2002

Data Year	County	24-Hour Lead			Qtr 1	Qtr 2	Qtr 3	Qtr 4	# Exceed	Monitor Number	Site ID	Site Address	City
		# Obs	1st Max	2nd Max									
1999	Shoshone	59	0.14	0.13	0.04	0.03	0.05	0.04	0	1	160790006	Medical Clinic/204 Oregon	Kellogg
2000	Shoshone	61	0.49	0.10	0.04	0.04	0.08	0.04	0	1	160790006	Medical Clinic/204 Oregon	Kellogg
2001	Shoshone	58	0.06	0.04	0.03	0.03	0.03	0.03	0	1	160790006	Medical Clinic/204 Oregon	Kellogg
2002	Shoshone	30	0.05	0.04	0.03	0.03			0	1	160790006	Medical Clinic/204 Oregon	Kellogg